

09674377 103000

09/674377

LAW OFFICES

SUGHRUE, MION, ZINN, MACPEAK & SEAS, PLLC

2100 PENNSYLVANIA AVENUE, N.W.

WASHINGTON, DC 20037-3213

TELEPHONE (202) 293-7060

FACSIMILE (202) 293-7860

www.sughrue.com

534 Rec'd PCT/PTG 30 OCT 2000

October 30, 2000

BOX PCTAssistant Commissioner for Patents
Washington, D.C. 20231

PCT/PCT/JP99/01834

-filed April 6, 1999

Re: Application of Toshikazu NAKAMURA
NEOVASCULARIZATION INHIBITORS
Our Ref: Q61434

Dear Sir:

The following documents and fees are submitted herewith in connection with the above application for the purpose of entering the National stage under 35 U.S.C. § 371 and in accordance with Chapter II of the Patent Cooperation Treaty:

- ☒ an executed Declaration and Power of Attorney.
- ☒ an English translation of the International Application.
- ☒ thirteen (13) sheets of drawings.
- ☒ eight (8) sheets of Sequence listing and a 3.5" disk containing Sequence listing in a computer readable form.
- ☒ Statement in Support of Submission in Accordance with 37 C.F.R. § 1.821.
- ☒ International Preliminary Examination Report.
- ☒ International Search Report and a Form PTO-1449 listing the ISR references.
- ☒ Translation of PCT Article 34 Amendment

It is assumed that copies of the International Application, the International Search Report, the International Preliminary Examination Report, and any Articles 19 and 34 amendments as required by § 371(c) will be supplied directly by the International Bureau, but if further copies are needed, the undersigned can easily provide them upon request.

The Government filing fee is calculated as follows:

Total claims	26	-	20	=	6	x	\$18.00	=	\$108.00
Independent claims	10	-	3	=	7	x	\$80.00	=	\$560.00
Base Fee									\$860.00
Multiple Dependent Claim Fee									\$270.00
TOTAL FILING FEE									\$1,798.00

The Office is hereby directed and authorized to charge the statutory filing fee of \$1180.00 to Deposit Account No. 19-4880. You are also directed and authorized to charge or credit any difference or overpayment to said Account. The Commissioner is hereby authorized to charge any fees under 37 C.F. §§ 1.16, 1.17 and 1.492 which may be required during the entire pendency of the application to Deposit Account No. 19-4880. A duplicate copy of this transmittal letter is attached.

Priority is claimed from April 28, 1998 based on Japanese Application No. 1998-134681.

Since October 28, 2000 (30 months from the priority date) fell on a Saturday, the submission of these papers on Monday, October 30, 2000, is sufficient for entry of National Stage of the above application.

Respectfully submitted,

Waddell A. Biggart
Waddell A. Biggart
Registration No. 24,861

Bruce E. Kramer
Reg. No. 33,725

WAB/amt

09674377 103000

09/674377

534 Rec'd PCT/PTO 30 OCT 2000

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Toshikazu NAKAMURA et al.

Appln. No.: unassigned

Group Art Unit: unassigned

Filed: October 30, 2000

Examiner: unassigned

For: NEOVASCULARIZATION INHIBITORS

**STATEMENT TO SUPPORT FILING AND SUBMISSION IN
ACCORDANCE WITH 37 C.F.R. §§ 1.821-1.825**

Assistant Commissioner for Patents

Washington, D.C. 20231

Box SEQUENCE

Sir:

In connection with a Sequence Listing submitted concurrently herewith, the undersigned hereby states that:

1. the submission, filed herewith in accordance with 37 C.F.R. § 1.821(g), does not include any new matter;
2. the content of the attached paper copy and the attached computer readable copy of the Sequence Listing, submitted in accordance with 37 C.F.R. § 1.821(c) and (e), respectively, are the same; and
3. all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true, and further, that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that

such willful false statements may jeopardize the validity of the application or any patent resulting therefrom.

Respectfully submitted,



Mark A. Hisson
Registration No. 44,765

SUGHRUE, MION, ZINN,
MACPEAK & SEAS, PLLC
2100 Pennsylvania Avenue, N.W.
Washington, D.C. 20037-3213
Telephone: (202) 293-7060
Facsimile: (202) 293-7860

Date: October 30, 2000

13/PRTS

09674377 103000

09/674377

534 Rec'd PCT/PTO 30 OCT 2000

1

DESCRIPTION

NEOVASCULARIZATION INHIBITORS

TECHNICAL FIELD

The present invention relates to neovascularization inhibitors. More particularly, the invention relates to a neovascularization inhibitor (antineovascularization composition) which comprises a protein having a defined regional sequence of the α -chain of hepatocyte growth factor (hereinafter referred to sometimes briefly as HGF) as an active ingredient.

The neovascularization inhibitor of the present invention finds application, based on its inhibitory effect on neovascularization, as a prophylactic or therapeutic agent for various diseases associated with abnormal angiogenesis, such as rheumatoid arthritis, diabetic retinopathy, retinopathy of prematurity, senile macular degeneration and hypercicatrization in wound healing.

BACKGROUND ART

Neovascularization is a phenomenon such that

the vascular endothelial cells, mainly of venules, form a de novo vasculature in response to some stimulus or other. In the normal state of a living body, this phenomenon is indispensable for sustained metabolism of tissues and for functional homeostasis of the body and is generally observed in the process of wound healing, growth of fetal lungs, and evolution of luteinization.

Meanwhile, it is well known that an abnormal angiopoiesis is involved in various diseases inclusive of inflammatory diseases. For example, such diseases as proliferating diabetes, psoriasis vulgaris, rheumatoid arthritis, diabetic retinopathy, senile macular degeneration, overcircatrization in wound healing, etc. and the metastasis and recurrence of carcinomas are reportedly caused by hyperplasia of blood vessels, particularly peripheral capillary vessels [Polverini PJ., Crit. Rev. Oral. Biol. Med., 1995, 6(3), pp.230-247, Review: Forkman J., Nature Med., 1995, 1(1), pp.27-31].

Therefore, as prophylactic or therapeutic drugs for those diseases, a variety of antineovascularization drugs comprising compounds having neovascularization-inhibitory activity as

active ingredients have been developed to this day.

The present invention has for its object to provide a novel antineovascularization factor. Another object of the present invention is to provide a neovascularization inhibitor useful for the prevention and treatment of said various diseases arising from hyperplasia of blood vessels.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 A is a reversed phase high-performance liquid chromatogram (reversed phase HPLC) (C4) of elastase-treated HGF and B is an electrophoretogram (under reducing and non-reducing conditions) of peak fractions on said reversed phase HPLC.

Fig. 2 is a schematic representation of the structures of the α - and β -chains of HGF and the structure of the PyrGlu³² - Val⁴⁷⁸ region of HGF (HGF/NK4) which has been excised by elastase treatment.

Fig. 3 A and B show the dose-dependent rates of binding of ¹²⁵I-HGF and ¹²⁵I-HGF/NK4, respectively, to the rat liver plasma membrane. The insets in Fig. 3 A and B show the rates of binding of ¹²⁵I-HGF and ¹²⁵I-HGF/NK4 to said rat liver plasma membrane in Scatchard plots. Fig. 3 C is a diagram showing the

rates of binding of unlabeled HGF and unlabeled HGF/NK4 to the rat liver plasma membrane in the presence of ^{125}I -HGF (cf. Example 2).

Fig. 4 is a diagram showing the presence or absence of mitogenic activity in HGF and HGF/NK4 (A) and a diagram showing the antagonistic action of HGF/NK4 against the mitogenic activity of HGF. The mitogenic activity information was generated by assaying the DNA synthesis of rat primary-culture hepatocytes. Specifically, Fig. 4 A shows the DNA synthesis of hepatocytes in the presence of HGF or HGF/NK4 and Fig. 4 B shows the effect of HGF/NK4 on the DNA synthesis of hepatocytes in the presence of 60 pM HGF or 1.5 nM epidermal growth factor (EGF) [cf. Example 3].

Fig. 5 is a diagram showing the inhibitory effect of HGF/NK4 on the proliferation of human lung microvascular endothelial cells in the presence or absence (None) of bFGF, HGF or VEGF [Example 7].

Fig. 6 is a diagram showing the inhibitory effect of HGF/NK4 on the proliferation of human skin microvascular endothelial cells in the presence or absence (None) of bFGF, HGF or VEGF [Example 7].

Fig. 7 is a diagram showing the effects of HGF/NK4 (NK4 on the drawing) and anti-HGF antibody

(α -HGF Ab on the drawing) on the proliferation of human capillary vessel endothelial cells in the presence of 5% fetal bovine serum (FBS), 5% FBS+bFGF, 5% FBS+VEGF or 5% FBS+HGF [cf. Example 9 A].

Fig. 8 is a diagram showing the effects of HGF/NK4 (NK4 on the drawing) and anti-HGF antibody (α -HGF Ab on the drawing) on the migration of human capillary vessel endothelial cells in the presence of 1% fetal bovine serum (FBS), 1% FBS+bFGF, 1% FBS+VEGF or 1% FBS+HGF [cf. Example 9 B].

Fig. 9 is a set of photographs, in lieu of a drawing, which shows the results of observation of the neovascularization-inhibitory effect of HGF/NK4 on chick embryonic chorioallantoic membrane with a stereoscopic microscope.

Fig. 10 is a photograph, in lieu of a drawing, which shows the inhibitory effect of HGF/NK4 on tumor neovascularization as assayed by an immuno histochemical method.

Fig. 11 indicates that HGF/NK4 has the action to inhibit growth of Lewis lung tumor cells, wherein A is a diagram showing the time course of the volume of a transplanted tumor mass and B is a histogram showing the weight of the transplanted tumor at day 28.

Fig. 12 A is a histogram indicating that HGF/NK4 has the action to inhibit metastasis of Lewis lung tumor cells [Reference Example 1] and Fig. 12 B is a photograph, in lieu of a drawing, which shows the metastasis of the tumor to the lung, indicating clearly that whereas a plurality of lung metastatic foci are present in the saline-treated control group, substantially no lung metastatic foci are detected in the HGF/NK4-treated group.

Fig. 13 indicates that HGF/NK4 has the action to inhibit the growth (A) and metastasis (B) of Jyg mammary tumor cells; A shows the time course of the volume of a transplanted tumor mass and B shows the number of metastatic foci.

DISCLOSURE OF INVENTION

Searching for a novel neovascularization inhibitor in earnest with the above object in mind, the inventors of the present invention found that a protein containing a defined region of the α -chain of hepatocyte growth factor (HGF) has the action to significantly inhibit neovascularization and have accordingly developed the present invention.

HGF is the very polypeptide which the inventors

discovered in 1984 as a novel growth factor for hepatic parenchymal cells (Nakamura, T. et al., Biochem Biophys Res Commun., 1984, 122, pp.1450-1459). Subsequent studies by the present inventors revealed that HGF is a heterodimer consisting of an α -chain having a molecular mass of about 69 kDa and a β -chain having a molecular mass of about 34 kDa and has a unique domain structure comprising an N-terminal hairpin domain and 4 Kringle domains in the α -chain and a serine protease-like domain in the β -chain (Nakamura T., et al., Nature 1989, 342, pp.440-443). Though, as of the time of its discovery, HGF was considered to be a growth factor with highly specificity to hepatocytes as the substantive entity of liver regeneration factor, the subsequent studies made since 1989 when recombinant HGFs became available revealed that HGF acts also as a potent mitogen for many kinds of epithelial cells in addition to hepatocytes (Nakamura T., Princess Takamatsu Symp., 1994, 24, pp.195-213. Review). Moreover, the results of further investigations indicated that, in addition to the above cell growth-regulating function, HGF not only has the function of a motogen enhancing cell motility (T. Nakamura, Prog. growth Factor Res., 3,

pp.67-85, 1991) but also has novel biological activities inclusive of tumor suppressant activity which inhibits proliferation of many kinds of tumor cells (Higashio K, et al., Biochem Biophys Res Commun., 1990, July 16, 170(1), pp.397-404).

Furthermore, it was elucidated, in 1991, that the functional receptor having a high affinity for HGF is a protooncogene product (c-met product: c-Met) (Bottaro D. P., et al., Science, 1991, Feb., 15, 251(4995), pp.802-804; Naldini L, et al., Oncogene. 1991 Apr, 6(4), pp.501-504), and the present inventors discovered that the N-terminal hairpin domain and first and second Kringle domains of the α -chain are the minimum domains binding to said c-Met/HGF receptor (Matsumoto, K. et al., Biochem. Biophys. Res. Commun., 1991 Dec., 16, 181(2), pp.691-699).

The present inventors did a further study on the heels of the above series of research into HGF and arrived at the novel finding that a polypeptide having said N-terminal hairpin and first through fourth Kringle domains of the α -chain has antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and confirmed that said polypeptide has the action to significantly

suppress the neovascularization-inducing action of HGF.

The present invention has been developed on the basis of years of those studies on HGF.

The present invention, therefore, is directed to the following neovascularization inhibitors 1-6.

1. A neovascularization inhibitor comprising the following polypeptide (a) or (b) as an active ingredient.

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor.

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence of (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

2. A neovascularization inhibitor comprising the following polypeptide (a) or (b) as an active ingredient:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or

AH34

more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF.

3. A neovascularization inhibitor as set forth in paragraph 1 or 2, wherein said polypeptide has at least one hairpin domain and 4 Kringle domains.

4. A neovascularization inhibitor as set forth in paragraph 1 or 2, wherein said polypeptide is one obtainable by elastase digestion of hepatocyte growth factor.

5. A neovascularization inhibitor comprising the polypeptide defined by SEQ ID NO:1 and a pharmaceutically acceptable carrier.

6. A neovascularization inhibitor comprising the polypeptide defined by SEQ ID NO:2 and a pharmaceutically acceptable carrier.

The present invention is further directed to the following medically or pharmacologically useful agents 7-11.

7. A prophylactic or therapeutic drug for a disease associated with abnormal angiopoiesis which comprises the polypeptide as set forth in paragraph 1 or 2 and a pharmaceutically acceptable carrier.

8. A prophylactic or therapeutic drug as set forth in paragraph 7 wherein said disease associated with abnormal angiopoiesis is selected from the group consisting of rheumatoid arthritis, psoriasis, Osler-Webber syndrome, myocardial angiopoiesis, telangiectasia, hemophilic joint, angiogenic diseases of the eye, angiofibroma, benign tumors and wound granulation.

9. A prophylactic or therapeutic drug for a disease arising from overstimulation of endothelial cells which comprises the polypeptide as set forth in paragraph 1 or 2 and a pharmaceutically acceptable carrier.

10. A prophylactic or therapeutic drug as set forth in paragraph 9 wherein said disease arising from overstimulation of endothelial cells is selected from the group consisting of enteric adhesion, Crohn's disease, atherosclerosis, scleroderma and overcicatrizization.

11. A conception-regulating drug comprising the polypeptide as set forth in paragraph 1 or 2 and a pharmaceutically acceptable carrier.

The present invention is further directed to the following medically or pharmacologically useful treatment methods 12-16.

H/31

12. A method of inhibiting neovascularization which comprises administering to a subject a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

13. A method of inhibiting neovascularization which comprises administering to a subject a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of

HKT34

13/1

HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF.

14. A method for prophylaxis or therapy of a disease associated with abnormal angiopoiesis which comprises administering a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF to a subject in whom a prophylactic or therapeutic treatment for said disease is indicated.

15. A method for prophylaxis or therapy of a disease associated with abnormal angiopoiesis which comprises administering a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

13/2

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF to a subject in whom a prophylactic or therapeutic treatment for said disease is indicated.

16. The method for prophylaxis or therapy as set forth in paragraph 14 or 15 wherein said disease is any disease selected from the group consisting of rheumatoid arthritis, psoriasis, Osler-Webber syndrome, myocardial angiopoiesis, telangiectasia, hemophilic joint, angiogenic diseases of the eye, angiofibroma, benign tumors, wound granulation, enteric adhesion, Crohn's disease, atherosclerosis, scleroderma and overcicatrizization.

In a further aspect, the present invention is directed to the use of the polypeptide 17-18.

17. Use of the following polypeptide (a) or (b) for the production of a neovascularization inhibitor:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

13/3

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

18. Use of the following polypeptide (a) or (b) for the production of a neovascularization inhibitor:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF.

Representation of amino acids, peptides, nucleotide sequences and others by abbreviations in this specification is principally in conformity with the nomenclature recommended by IUPAC and IUPAC-IUB and the rules set forth in the "Guideline for Preparation of a Specification or the Equivalent Referring to a Nucleotide Sequence or an Amino Acid

ART 34

13/4

Sequence" (March 1997, The Examination Standards Office, Coordination Division, the Patent Office of Japan).

Furthermore, the amino acid numbers and positions as mentioned in this specification are based on the amino acid sequence of prepro-HGF (Nakamura T., et al., Nature 1989, 342, pp.440-443).

ART 34

14/1

In the context of this invention, PyrGlu means pyroglutamate, which is a modified amino acid residue, and PyrGlu³² signifies that, based on the amino acid sequence of prepro-HGF, the 32nd amino acid residue from the N-terminus is pyroglutamate.

The neovascularization inhibitor (antineovascularization composition) of the present invention contains as an active ingredient a polypeptide resulting from the α -chain of HGF and having neovascularization inhibitory activity.

The term "resulting from the α -chain of HGF" is used herein to mean that the entire region or fragments of the α -chain of HGF are contained, whether continuously or discontinuously.

The term neovascularization inhibitory activity means any action that suppresses neovascularization without regard to its mode or mechanism. In a preferred sense, it means the action to suppress the neovascularization-inducing action of HGF. More preferably, it means the action to suppress the neovascularization-inducing action of HGF, VEGF or bFGF.

As polypeptides having such activity, there can be mentioned polypeptides having an affinity for the c-Met/HGF receptor, which is the receptor of HGF,

HAY 3U

14/2

and the action to competitively antagonize the binding of HGF to said receptor. The preferred are

polypeptides which bind to the c-Met/HGF receptor to exhibit antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

The c-Met/HGF receptor-mediated action of HGF includes c-Met/HGF receptor tyrosine phosphorylating activity, motogenic activity, mitogenic activity and morphogenic activity [Jikken Igaku (Experimental Medicine), Vol. 11, No. 9 (1993)].

Therefore, the polypeptide for use in the present invention is a polypeptide which suppresses or inhibits any of such activities, more particularly a polypeptide having at least one action selected from among the action to suppress/inhibit the HGF-induced c-Met/HGF receptor tyrosine phosphorylation, the action to suppress/inhibit the motogenic activity of HGF, the action to suppress/inhibit the mitogenic activity of HGF and the action to suppress/inhibit the morphogenic activity of HGF. Preferably, a polypeptide having all of the above-mentioned actions can be mentioned.

As preferred examples of such polypeptide, there can be mentioned a polypeptide having the amino acid sequence $\text{PyrGlu}^{32} \sim \text{Val}^{478}$ of HGF, more

particularly a polypeptide having the amino acid sequence depicted in SEQ ID NO:1. In SEQ ID NO:1, "Xaa" represents "PyrGlu".

This polypeptide can be obtained basically by subjecting HGF to elastase treatment, and consists of the 447 amino acids in the N-terminal region of the α -chain of HGF. Therefore, like the α -chain of HGF, this particular polypeptide has the modified amino acid residue = pyroglutamate at the N-terminus and contains one N-terminal hairpin domain and 4 Kringle domains. In this specification, the above polypeptide is referred to sometimes as HGF/NK4.

As will be shown in the Example which appears hereinafter, this polypeptide exhibits said antagonistic activity by binding to the c-Met/HGF receptor in competition with HGF and has the action to suppress or inhibit the HGF-induced autotirosine-phosphorylation of the c-Met/HGF receptor. Furthermore, this polypeptide has little mitogenic, motogenic or morphogenic activities of its own and has the property to inhibit the mitogenic, motogenic and morphogenic activities of HGF.

However, the polypeptide for use as the active ingredient of the neovascularization inhibitor of the present invention is a polypeptide resulting

from the α -chain of HGF and, inasmuch as it has the action to inhibit the neovascularization-inducing action of HGF, is not limited to said particular polypeptide shown in SEQ ID NO:1.

As specific examples of said polypeptide, there can be mentioned polypeptides having amino acid sequences derived from the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of HGF by the deletion, substitution or addition of one, a few or more amino acids and yet having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

More preferably, polypeptides equivalent or comparable to said HGF/NK4 in physiological activities (tyrosine phosphorylation inhibiting activity, anti-motogenic activity, anti-mitogenic activity, and anti-morphogenic activity) can be mentioned.

The extent of said "deletion, substitution or addition" of amino acids and the positions involved are not particularly restricted only if the mutant polypeptide still retains the above-mentioned physiological activities. By way of example, a polypeptide resulting from the deletion or addition of one or more than one (or several) amino acids in the N-terminal and/or C-terminal region of said

HGF/NK4 and a polypeptide resulting from the deletion or addition of one or more than one (or several) amino acids in the intermediate region of HGF/NK4. Preferably, however, at least one hairpin domain and 4 Kringle domains, which characterize the structure of HGF/NK4, are substantially retained after the mutation.

As a corollary, as a typical mutant peptide, a polypeptide resulting from the substitution, deletion or addition of one or more than one (or several) of the amino acids in the region exclusive of said hairpin domain and 4 Kringle domain can be mentioned. As specific examples of such mutant peptide, there can be mentioned a polypeptide [HGF/NK4 (del 5)] resulting from the deletion of 5 amino acids, namely amino acid Nos. 162~166 (amino acids Nos. 131~135 in SEQ ID NO:1), from the HGF/NK4 polypeptide, that is to say the polypeptide having the amino acid sequence depicted in SEQ ID NO:2.

HGF/NK4 or HGF/NK4 (del 5) for use in the present invention can be respectively produced chemically by a routine method for peptide synthesis or by a routine genetic engineering technique, on the basis of the amino acid sequence information given in SEQ ID NO:1 or 2 or the already-known gene

(nucleotide) sequence of HGF.

Preferably, it can be obtained by enzymatic degradation of HGF.

The enzymatic degradation of HGF can be achieved by digesting HGF with an elastase or the like enzyme. Then, this enzymatic digest is purified by the conventional protein purification method, for example high-performance liquid chromatography or SDS-PAGE, and a polypeptide having a given molecular mass is isolated to acquire HGF/NK4. The molecular mass mentioned above may be about 65~69 kD, preferably about 67 kD, as determined by SDS-PAGE under reducing conditions and about 48~52 kD, preferably about 50 kD, as determined by SDS-PAGE under non-reducing conditions.

HGF for use in said enzymatic degradation is not particularly limited in terms of its source or preparation procedure.

For example, it can be obtained by extraction and purification from liver or other tissues, blood cells such as platelets, leukocytes, etc., plasma or serum of mammals inclusive of man (FEBS, 224, 312, (1987); Proc. Acad. Sci. USA, 86, 5844 (1989)) or by growing primary-culture HGF-producing cells or

cell lines and separating and purifying HGF from the resulting culture. As a further alternative, a recombinant HGF can be acquired by a genetic engineering technique which may comprise cloning an HGF-encoding gene in a suitable vector, introducing the vector into suitable host cells (e.g. animal cells) for transfection and harvesting the objective recombinant HGF from a culture supernatant of the resulting cells (e.g. Nature, 342, 440, 1989; JP 1993-111383A; JP 1991-255096A; Biochem., Biophys. Res. Commun., 163, 967, 1989, etc.). The HGF-encoding gene which can be used routinely includes HGF genes derived from mammals inclusive of humans, preferably the HGF gene of the human origin, more preferably human-derived recombinant HGF genes (JP 1993-111383A).

The so-called mutants constructed on the basis of the amino acid sequence of HGF/NK4, such as HGF/NK4 (del 5), can be prepared chemically by a method for peptide synthesis or by a genetic engineering technique starting with an HGF gene.

The technology for mutagenesis includes genetic engineering methods such as site-specific mutagenesis [Methods in Enzymology, 154: 350, pp.367-382 (1987); ditto 100: 468 (1983); Nucleic

Acids Res., 12: 9441 (1984); Zoku Seikagaku Jikken Koza 1 "Idenshi Kenkyuho II" (Experimental Biochemistry Series 1 "Methods for Gene Research II") (edited by Japanese Biochemical Society), p105 (1986)], as well as the methods of chemical synthesis, such as the phosphotriester method and phosphoamidate method [J. Am. Chem. Soc., 89: 4801 (1967); do: 91: 3350 (1969); Science, 150: 178 (1968); Tetrahedron Lett., 22: 1859 (1981); do 24: 245 (1983)] and any combination of such techniques.

The neovascularization inhibitor of the present invention is not particularly restricted in dosage form and may be provided in a variety of dosage forms, namely oral preparations such as powders, fine granules, granules, tablets, pills, capsules, solutions, emulsions, suspensions, syrups, etc.; preparations for external application, such as ointments, creams, DDS patches, suppositories, etc.; ophthalmic preparations such as eyedrops, ophthalmic ointments, etc.; injections and drip infusions. These dosage forms can be manufactured by the pharmaceutical procedures well established in the art.

Regarding the manufacture of tablets, the carrier that can be used includes various excipients

such as lactose, sucrose, sodium chloride, glucose, urea, starch, calcium carbonate, kaolin, crystalline cellulose, silica, etc.; binders such as simple syrup, glucose solution, starch solution, gelatin solution, carboxymethylcellulose, shellac, methylcellulose, potassium phosphate, polyvinylpyrrolidone, etc.; disintegrators such as dried starch, sodium alginate, agar powder, laminaran powder, sodium hydrogencarbonate, calcium carbonate, polyethoxylated sorbitan fatty acid esters, sodium lauryl sulfate, stearyl monoglyceride, starch, lactose, etc.; disintegration inhibitors such as sucrose, stearic acid, cacao butter, hydrogenated oil, etc.; absorption promoters such as quaternary ammonium bases, sodium lauryl sulfate, etc.; humectants such as glycerin, starch, etc.; adsorbents such as starch, lactose, kaolin, bentonite, colloidal silica, etc.; and lubricants such as purified talk, salts of stearic acid, boric acid powder, polyethylene glycol and so on. Furthermore, where necessary, tablets may be manufactured in the form of coated tablets, i.e. tablets carrying a conventional surface coating, such as sugar-coated tablets, gelatin-coated tablets, film-coated tablets, etc.,

or in the form of double-layered or multi-layered tablets.

The carrier which can be used in the manufacture of pills include but is not limited to various excipients such as glucose, lactose, starch, cacao butter, hydrogenated vegetable oils, kaolin, talc, etc.; binders such as gum arabic powder, tragacanth powder, gelatin, etc.; and disintegrators such as laminaran and agar, among others.

Capsules can be manufactured by the conventional method which comprises blending said peptide with various carrier substances such as those mentioned above and filling the mixture into hard gelatin capsule shells, soft gelatin capsule shells or the like.

For use in the manufacture of suppositories, the carrier includes polyethylene glycol, cacao butter, higher alcohols, esters of higher alcohols, gelatin and semisynthetic glycerides, among others.

Injections can be manufactured by the conventional technology, for example by dissolving said polypeptide in a suitable solvent, sterilizing the solution for example by filtration, and distributing it into sterile vials. Such injections are preferably isotonic to blood, and the

diluent which can be used for provision of such dosage forms includes but is not limited to sterile water, ethyl alcohol, macrogols, propylene glycol, ethoxylated isostearyl alcohol, polyoxylated isostearyl alcohol, polyethoxylated sorbitan fatty acid esters. A sufficient amount of sodium chloride, glucose or glycerin to isotinize such injections may be included in formulations and the conventional solubilizer, buffer, local anesthetic and other additives may also be added.

For the manufacture of an ointment, the ointment base, stabilizer, lubricant, preservative, etc., which are usually employed for such peptides, are formulated and processed in the conventional manner to provide the objective product. The ointment base mentioned above includes liquid paraffin, white petrolatum, bleached beeswax, paraffin and so on. The preservative includes methyl p-hydroxybenzoate, ethyl p-hydroxybenzoate and propyl p-hydroxybenzoate, among others.

Transdermal drug delivery systems can be prepared by spreading said ointment or an equivalent thereof in the form of a paste, cream or gel on the conventional support in the conventional manner. The suitable support includes a woven or nonwoven

cloth of cotton, spun rayon or other chemical fiber or a flexible polyvinyl chloride, polyethylene, polyurethane or other sheet or foam sheet.

Furthermore, each of said various pharmaceutical preparations may be supplemented, where necessary, with various pharmaceutically acceptable additives such as a coloring agent, a preservative, a flavoring agent, a corrigent, a sweetener, etc. and/or other medicaments.

In pharmaceutical production runs, a stabilizer is preferably formulated. The stabilizer which can be used includes albumin, globulin, gelatin, mannitol, glucose, dextran and ethylene glycol, to mention but a few examples.

Solutions inclusive of parenteral preparations should be stored frozen or preferably supplied as lyophilized products. Lyophilized preparations are extemporaneously reconstituted with distilled water for injection or the like solvent vehicle.

The amount of the polypeptide to be formulated in the pharmaceutical composition of the present invention is not particularly restricted but can be liberally selected from a broad range. Generally, however, the recommended concentration in the

composition is about 0.0002~0.2 (w/v) %, preferably about 0.001~0.1 (w/v) %.

The method of administering the pharmaceutical composition is not particularly restricted but can be judiciously selected with reference to the dosage form, patient factors such as age, sex, etc. and severity of illness, among other conditions. By way of illustration, parenteral preparations may be intravenously administered either independently or in admixture with the conventional glucose, amino acid or other infusion. Where necessary, such preparations may be administered alone intramuscularly, intradermally, subcutaneously or intraperitoneally.

The daily dosage of the neovascularization inhibitor of the present invention is dependent on the patient's condition, body weight, age, sex and other factors and cannot be stated in general terms. However, in terms of the amount of the polypeptide of the invention (HGF/NK4 or a mutein thereof), the recommended usual dosage for an adult human is about 0.01~100 mg/day and this dose may be administered in a single dose or in a few divided doses.

The neovascularization inhibitor of the present invention can be applied to the prophylaxis

and therapy of a broad spectrum of diseases arising from vascular hyperplasia. Although such diseases are not particularly restricted, there can be mentioned rheumatoid arthritis, psoriasis, Osler-Webber syndrome, myocardial angiopoiesis, telangiectasia, hemophilic joint, angiogenic diseases of the eye (e.g. diabetic retinopathy, retinopathy of prematurity, senile macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, etc.), angiofibroma, benign tumors (e.g. hemangioma, acoustic neuroma, neurofibroma, trachoma, granuloma pyogenicum, etc.), hematopoietic malignancies such as leukemia, solid cancers, cancer metastases and wound granulation, among others.

The neovascularization inhibitor of the present invention is further applicable broadly to the prophylaxis and therapy of diseases caused by an excessive or abnormal stimulation of the endothelial cells. Such diseases are not particularly restricted but include, among others, enteric adhesion, Crohn's disease, atherosclerosis, scleroderma, and overcicatrizization such as keloid.

Furthermore, the neovascularization inhibitor

of the present invention finds application as a drug for conception control based on inhibition of the neovascularization necessary for implantation and finds application as a prophylactic or therapeutic drug for diseases accompanied by angiogenesis, such as cat scratch disease and ulcer, as an pathologic outcome.

BEST MODE FOR CARRYING OUT THE INVENTION

The following examples are intended to illustrate the present invention in further detail without delineating the technical scope of the invention. Thus, many modifications and changes can be made by those skilled in the art based on the disclosure in this specification without departing from the technical scope of the invention.

Example 1 Isolation and purification of HGF/NK4

CHO cells transfected with human HGF cDNA were cultured and the recombinant HGF was isolated and purified from the culture (Nature 342, pp.440-443 (1989); Biochem. Biophys. Res. Commun. 172; pp.321-327 (1990)). This recombinant HGF (900 mg) was digested with pancreatic elastase (Sigma) in 0.2 M Tris-HCl buffer (pH 8.8) at 37°C for 20 minutes (enzyme:substrate = 1:100). This digest was

purified by reversed-phase high-performance liquid chromatography (HPLC; μ Bondapak C4 column, Waters Japan) on a gradient of 0.05% trifluoroacetic acid-acetonitrile. On the HPLC, three peaks were detected as shown in Fig. 1A.

Then, each of these peak fractions was subjected to SAS-PAGE and protein staining (Fig. 1B). It was found that a first peak is a polypeptide fragment having a molecular mass of 50 kD under nonreducing conditions and of 67 kD under reducing conditions; a second peak corresponds to the undigested heterodimer HGF consisting of an α -chain of 69 kD and a β -chain of 34/32 kD; and a third peak is a fragment having a molecular mass of 33/31 kD under nonreducing conditions and of 34/32 kD under reducing conditions.

In the above description, the expression $_/_$ kD indicates the existence of proteins of the same amino acid sequence but different molecular masses depending on differences in the appended sugar chain.

The foregoing indicated that, upon elastase treatment, HGF is digested into two fragments, namely a fragment (the first peak) comprising a major part of the α -chain but being slightly smaller

than the full-length α -chain and a fragment (the third peak) consisting of part of the C-terminal sequence of the α -chain and the full-length β -chain.

For the amino acid sequencing of the purified fragments, the solvent was first evaporated off from each fraction of the eluate and the residue was dissolved in 0.1 M phosphate buffer (pH 7.3) containing 0.05% CHAPS (Sigma) and 1 M NaCl. The amino acid analysis was carried out using Automatic Protein Sequencer 492 (Applied Biosystem Inc.).

To begin with, the analysis for the N-terminal sequence of the first-peak fragment was attempted but failed, suggesting the likelihood that the N-terminus of this fragment was in the pyroglutamate form as it was the case with the N-terminus of HGF. Therefore, this fragment was first treated with pyroglutamate aminopeptidase and then analyzed for the N-terminal sequence. As a result, the sequence, as expressed in single capital letters, was found to be RKRRNTIHEF in agreement with the N-terminal amino acid sequence No. 2 ~ No. 11 of the α -chain. It was thus found that the N-terminal amino acid of this fragment was glutamine modified in the pyroglutamate form and the structure of the N-

terminal region was identical with that of undigested HGF.

Then, for analysis of the C-terminus of this fragment, an amino acid analysis was carried out on the other fragment produced by elastase digestion (the third fragment) for the N-terminus of its partial α -chain. As a result, the amino acid sequence of the N-terminal region of this partial α -chain was found to correspond to the Asn⁴⁷⁹ ~ Ala⁴⁸⁸ of HGF. This finding indicated that the C-terminus of the first-peak fragment was Val⁴⁷⁸.

Based on the above results, it was clear that HGF, on elastase treatment, is digested into two fragments, one of which is a fragment consisting in the PyrGlu³² ~ Val⁴⁷⁸ region having the hairpin domain and 4 Kringle domains of HGF (HGF/NK4) while the other is a fragment consisting of part of the α -chain (16 amino acids starting with Asn⁴⁷⁹) and the β -chain (Fig. 2).

Example 2 Binding affinity of HGF/NK4 for cell surface receptors

Using the HGF/NK4 prepared in Example 1, its binding affinity for the cell surface receptor was investigated. As a control experiment, the binding affinity of HGF for the cell surface receptor was

also investigated. In this connection, the HGF/NK4 and HGF were radio-labeled (^{125}I -HGF/NK4 and ^{125}I -HGF) by the chloramine-T method, and as the cell surface receptor sample, the plasma membrane prepared from the rat liver was used.

Scatchard analysis revealed that both HGF and HGF/NK4 bind to the cell surface receptors in a concentration-dependent manner up to 80 pM (Fig. 3 A and B), that the K_d and number of HGF receptors were 64.5 pM and 5478 sites/ng, respectively, and that the K_d and number of HGF/NK4 receptors were 486 pM and 6427 sites/ng, respectively.

Then, to investigate whether HGF/NK4 is a competitive antagonist of HGF, the liver plasma membrane (50 μg) was incubated in the presence of ^{125}I -HGF alone (60 pM) or a mixture of ^{125}I -HGF and a varying amount of unlabeled HGF or unlabeled HGF/NK4. The membrane binding of ^{125}I -HGF was completely inhibited by the addition of unlabeled HGF and the 50% inhibitory concentration of unlabeled HGF was 60 pM. Unlabeled HGF/NK4 also inhibited the membrane binding of ^{125}I -HGF and the 50% inhibitory concentration of this HGF/NK4 was 600 pM, with the membrane binding of ^{125}I -HGF being completely inhibited at 60 nM [Fig. 3 C].

These results suggested that HGF/NK4 has an affinity for the c-Met/HGF receptor, although it is 8~10 times as low as that of HGF and, thus, is an antagonist of HGF.

Example 3 Mitogenic activity of HGF/NK4 and the inhibitory effect of HGF/NK4 on the mitogenic activity of HGF

The mitogenic activity of HGF/NK4 was evaluated by assaying the DNA synthesis of rat primary-culture hepatocytes. As a control experiment, the mitogenic activity of HGF was similarly evaluated (Nature 342, 440-443 (1989); Biochem. Biophys. Res. Commun. 181, 691-699 (1991)).

The data are presented in Fig. 4 A. It can be seen from the diagram that whereas HGF promoted the DNA synthesis of hepatocytes dose-dependently, HGF/NK4 did not promote the DNA synthesis even at a high concentration of 100 nM.

On the other hand, when HGF/NK4 was added to an HGF-containing culture medium so that both HGF and HGF/NK4 would be present, HGF/NK4 dose-dependently inhibited the DNA synthesis promoted by HGF, causing a substantially complete inhibition at 60 nM (Fig. 4 B). In contrast, HGF/NK4 showed no

inhibitory effect on the DNA synthesis promoted by epidermal growth factor (EGF) (Fig. 4 B).

These results indicated that although HGF/NK4 has no mitogenic activity of its own, it has an action to specifically inhibit the mitogenic action of HGF.

Example 4 Motogenic activity of HGF/NK4 and the antagonistic action of HGF/NK4 against motogenic activity of HGF

Using MDCK cells, the renal tubule-derived normal epithelial cells, the motogenic activity of HGF/NK4 was evaluated.

The MDCK cells formed a confluent mass of colonies in an HGF-free control medium but addition of HGF (22 pM) to the medium resulted in increased motility of MDCK cells and dispersion of the cells. In contrast, HGF/NK4 did not disperse the cells, with the mutual contact of cells being well maintained. Moreover, when MDCK cells were cultured in the presence of both HGF and HGF/NK4, HGF/NK4 inhibited the HGF-induced cell dispersion.

These results indicated that although HGF/NK4 has no motogenic activity of its own, it has an action to inhibit the motogenic activity of HGF.

Example 5 Morphogenic activity of HGF/NK4 and the antagonistic action of HGF/NK4 against the

morphogenic activity of HGF

To explore whether HGF/NK4 inhibits the morphogenic activity of HGF, MDCK cells, the renal tubule-derived normal epithelial cells, were cultured in collagen gel in the presence of HGF and/or HGF/NK4.

While the MDCK cells formed spherical cell masses in an HGF-free control medium, addition of HGF (55 pM) to the medium induced formation of a branched luminal structure. Addition of HGF/NK4 (55 nM) to this system did not induce such a luminal structure. Moreover, when MDCK cells were grown in the presence of both HGF and HGF/NK4, the cells remained in the form of masses without induction of a luminal structure.

The above results indicated that although HGF/NK4 has no morphogenic activity of its own, it has an action to inhibit the morphogenic activity of HGF.

Example 6 Inhibitory effect of HGF/NK4 on HGF-induced c-Met tyrosine phosphorylation

The lung tumor cell line A549 is known to have c-Met/HGF receptors expressed and to show enhanced tyrosine phosphorylation of c-Met in response to HGF stimulation. Therefore, it was explored whether

HGF/NK4 would inhibit the tyrosine phosphorylation of c-Met by HGF.

Thus, A549 cells were solubilized by stimulation with HGF and/or HGF/NK4 and immunoprecipitated with anti-c-Met antibody. Western blotting was then carried out and the tyrosine phosphorylation of c-Met was studied using anti-phosphotyrosine antibody.

As a result, whereas the stimulation with HGF (110 pM) was found to induce the tyrosine phosphorylation of c-Met, the stimulation with HGF/NK4 (110 nM) induced little phosphorylation. Moreover, HGF/NK4 dose-dependently inhibited the HGF-induced tyrosine phosphorylation of c-Met.

The above results indicated that HGF/NK4 inhibits the HGF-induced tyrosine phosphorylation of c-Met/HGF receptors. It was also suspected that, based on this inhibitory action, HGF/NK4 antagonizes the biological activity of HGF.

Example 7 Inhibitory effect of HGF/NK4 on growth of vascular endothelial cells

Using human lung microvascular endothelial cells (HMVEC-L; Clonetics) and human dermal microvascular endothelial cells (HMVEC-D; Clonetics) as tester vascular endothelial cells,

the growth inhibitory effect of HGF/NK4 on endothelial cells was evaluated.

Thus, using human lung microvascular endothelial cells or human skin microvascular endothelial cells in the logarithmic phase of growth of passage 5-8, a cell suspension was prepared and a gelatin-coated 24-well plate was seeded with 8000 cells per well. After 24 hours, the medium was changed to the fresh one (a 1:1 mixture of EGM (Eagle General Medium) and DMEM (Dulbecco's Modified Eagle Medium) supplied with 5% serum), and four groups of 3 ng/ml bFGF (basic fibroblast growth factor), 10 ng/ml HGF, 10 ng/ml VEGF (vascular endothelial growth factor) and negative control (5% serum-containing solution; None on the drawing) were established. Then, HGF/NK4 in a varying concentration of 0 to 450 nM was added and the plate was incubated under 5% CO₂ at 37°C. After 72 hours, the cells were detached by trypsin coating and counted using a Coulter counter.

The results for human lung microvascular endothelial cells are shown in Fig. 5 and those for human skin microvascular endothelial cells are shown in Fig. 6.

It can be seen from these diagrams that HGF/NK4

AK134

inhibits growth of vascular endothelial cells as induced by stimulation with 3 ng/ml bFGF, 10 ng/ml HGF, 10 ng/ml VEGF and 5% serum, respectively, all concentration-dependently and significantly.

These results suggested that HGF/NK4 acts in an inhibitory way not only against HGF-induced growth of vascular endothelial cells but also against the growth induced by other vascular endothelial cell growth factors such as bFGF and VEGF.

Example 8 Effects of anti-HGF antibody and HGF/NK4 on human capillary vessel endothelial cells

A. Influence on cell growth

The effects of anti-HGF antibody and HGF/NK4 on growth of human capillary vessel endothelial cells were evaluated.

Method

Cultured human skin capillary vessel endothelial cells were washed with phosphate-buffered saline (PBS) and detached with trypsin-EDTA (phosphate-buffered saline (PBS) containing 0.05% trypsin and 0.02% EDTA). These endothelial cells were suspended in EBM-2 medium (Clonetics) supplemented with 5% fetal bovine serum (FBS), seeded on a gelatin-coated 24-well plate at a density of 5×10^3 cells/cm² and cultured for 24 hours.

The culture was divided into 10 groups according to the following 10 kinds of media based on 5% FBS-EBM-2 medium and the culture medium was changed to the respective media. The symbol α -HGF Ab which appears below stands for anti-HGF rabbit polyclonal antibody.

1. 5% FBS
2. 5% FBS + 3 ng/ml bFGF
3. 5% FBS + 3 ng/ml bFGF + 300 nM HGF/NK4
4. 5% FBS + 3 ng/ml bFGF + 10 μ g/ml α -HGF Ab
5. 5% FBS + 10 ng/ml VEGF
6. 5% FBS + 10 ng/ml VEGF + 300 nM HGF/NK4
7. 5% FBS + 10 ng/ml VEGF + 10 μ g/ml α -HGF Ab
8. 5% FBS + 3 ng/ml HGF
9. 5% FBS + 3 ng/ml HGF + 300 nM HGF/NK4
10. 5% FBS + 3 ng/ml HGF + 10 μ g/ml α -HGF Ab

These media were incubated at 37°C under 5% CO₂ for 72 hours, after which time the cells were detached by trypsin coating and counted with a Coulter counter.

Results

The results are shown in Fig. 7. It can be seen from the diagram that 300 nM HGF/NK4 definitely inhibited growth of vascular endothelial cells as promoted by stimulation with 3 ng/ml bFGF, 10 ng/ml

VEGF and 3 ng/ml HGF, respectively. On the other hand, 10 μ g/ml anti-HGF rabbit polyclonal antibody specifically inhibited the HGF-stimulated growth of vascular endothelial cells but did not inhibit the growth of vascular endothelial cells stimulated with bFGF or VEGF. These results suggested that, aside from the HGF-antagonizing action, HGF/NK4 inhibits growth of vascular endothelial cells through some other novel activity.

B. Influence on cell migration

The effects of anti-HGF antibody and HGF/NK4 on migration of human capillary vessel endothelial cells were evaluated.

Method

The effect on cell migration was studied by the Boiden chamber method (Yoshida, A. et al., Growth Factors, 1996; 13(1-2): 57-64). Thus, the Boiden chamber test was carried out using a polycarbonate filter with a pore size of 5 μ m which had been coated with 13.4 μ g/ml of fibronectin.

First, human capillary vessel endothelial cells were cultured in serum-free EBM-2 medium for 12 hours and suspended in EBM-2 containing 1% fetal bovine serum and the suspension was seeded on said filter at a density of 12×10^4 cells/cm².

The culture was divided into 10 groups, and bFGF, HGF, VEGF, HGF/NK4 and anti-HGF rabbit polyclonal antibody (α -HGF Ab) were respectively added to the external fluid of the filter cup in these groups as follows.

1. None (1% FBS-EBM-2 medium)
2. 3 ng/ml bFGF
3. 3 ng/ml bFGF + 300 nM HGF/NK4
4. 3 ng/ml bFGF + 10 μ g/ml α -HGF Ab
5. 10 ng/ml VEGF
6. 10 ng/ml VEGF + 300 nM HGF/NK4
7. 10 ng/ml VEGF + 10 μ g/ml α -HGF Ab
8. 3 ng/ml HGF
9. 3 ng/ml HGF + 300 nM HGF/NK4
10. 3 ng/ml HGF + 10 μ g/ml α -HGF Ab

These samples were incubated for 5 hours and the number of cells (per visual field) which had migrated to the underside of the filter was determined under the microscope ($\times 200$). For improved accuracy of determination, cell counting was performed in 5 randomly selected visual fields.

Results

The results are shown in Fig. 8. It is apparent from the diagram that 300 nM HGF/NK4 definitely inhibited the migration of vascular endothelial

cells stimulated by 3 ng/ml bFGF, 10 ng/ml VEGF or 3 ng/ml HGF. On the other hand, 10 μ g/ml anti-HGF rabbit polyclonal antibody specifically inhibited the HGF-stimulated migration of vascular endothelial cells but did not inhibit the bFGF or VEGF-stimulated migration of these cells. Those results suggested that, in addition to HGF-antagonizing activity, HGF/NK4 inhibits migration of vascular endothelial cells through some other novel activity.

Example 9 Inhibitory effect of HGF/NK4 on the neovascularization of chick chorioallantoic membrane (CAM)

Fertile chicken eggs were incubated for 4 days, after which the eggshell was drilled in two positions, namely over the air chamber and the lateral side of the shell. From the side hole, 3 ml of the egg white was aspirated and the shell was sealed with a tape. The shell and shell membrane over the air chamber were removed and a silicone ring was set centrally on the chorioallantoic membrane (CAM). Then, an HGF/NK4- or bovine serum albumin (control)-containing methylcellulose disk was set in the silicone ring. After 2 days of incubation at 37°C, the vasculature on the CAM was examined with

a stereoscopic microscope. The results are shown in Table 1.

Table 1

Degree of neovascular invasion	
HGF/NK4	+
Control	+++

The degree of neovascular invasion into the disk was evaluated on the following scale.

- : no invasion
- +: about 1-2 invading vessels
- ++: about 3-4 invading vessels
- +++: 5 or more invading vessels

The photographs of the findings obtained in the examination with a stereoscopic microscope, in lieu of a drawing, are presented in Fig. 9.

It will be apparent that whereas the control disk showed a marked neovascular invasion, the HGF/NK4 disk showed little evidence of neovascular invasion. This finding indicated that HGF/NK4 has neovascularization inhibitory activity.

Example 10 Inhibitory effect of HGF/NK4 on tumor neovascularization

Using 6-8-week-old nude mice (BALB/c nu/nu), 5×10^6 GB-d1 human gallbladder cancer cells were

transplanted subcutaneously in the dorsal region. After 7 days, an osmotic pressure pump (Alzet) containing HGF/NK4 or, as control, saline was implanted beneath the dorsal skin and HGF/NK4 or saline was continuously infused into the vicinity of the transplanted cancer cells for 13 days. At week 4 after transplantation, the tumor mass was excised, fixed and sectioned in the routine manner to prepare a tissue specimen. For evaluating neovascularization in the cancer tissue, the tissue section was stained by the immunohistochemical method using anti-von Willebrand factor antibody (Dako). The results are shown in Table 2.

Table 2

von Willebrand factor stain	
HGF/NK4	+
Control	+++

The degree of neovascularization in the cancer tissue was evaluated on the following scale.

- : no von Willebrand factor-positive microvessels
- + : about 1/4 of microvessels are von Willebrand factor-positive
- ++ : about 1/2 of microvessels are von Willebrand

factor-positive

+++: about 3/4 of microvessels are von Willebrand
factor-positive

++++: more than 3/4 of microvessels are von
Willebrand factor-positive

Moreover, the findings in the microscopic
examination are presented as photographs, in lieu
of a drawing [Fig. 10].

It will be apparent from the above results that
a large number of von Willebrand factor-positive
microvessels were found in the control saline-
infused graft tumor tissue, with no evidence of
apoptosis of cancer cells in the tumor mass. In
contrast, in the HGF/NK4-perfused graft tumor
tissue, tumor neovascularization was remarkably
inhibited, with evidence of an extensive apoptosis
of cancer cells in the center of the tumor mass.
These results indicated that HGF/NK4 inhibits tumor
neovascularization in vivo.

Reference Example 1 Inhibitory effect of HGF/NK4
on tumor growth and metastasis (1)

Using 6~8-week-old nude mice (BALB/c nu/nu),
 5×10^6 Lewis lung cancer cells were transplanted
subcutaneously at the animal back. After 5 days,
an osmotic pressure pump (Alzet) containing HGF/NK4

or saline (control) was implanted beneath the dorsal skin and HGF/NK4 or saline was continuously infused into the neighborhood of the tumor graft for 2 weeks. At 28 days after transplantation, the weight and pulmonary metastasis of the transplanted tumor were investigated. The tumor volume was calculated by means of the formula: $(\text{minor diameter})^2 \times (\text{major diameter})^2 \times 0.5$.

The time course of tumor volume is shown in Fig. 11 A and the weight of the transplanted tumor at day 28 is shown in Fig. 11 B. It will be apparent from these diagrams that whereas the transplanted tumor perfused with saline showed rapid growth at 10 days after transplantation and onwards, HGF/NK4 inhibited tumor growth strongly and dose-dependently.

Regarding the pulmonary metastasis of Lewis lung cancer, whereas a large number of pulmonary metastatic foci were observed in the control saline-perfused group, pulmonary metastasis was inhibited remarkably and dose-dependently in the animals perfused with HGF/NK4 (Fig. 12 A, B).

These results indicated that HGF/NK4 has an action to significantly inhibit the growth and metastasis of lung cancer in vivo.

Reference Example 2 Inhibitory effect of HGF/NK4 on cancer growth and metastasis (2)

Using 6~8-week-old nude mice (BALB/c nu/nu), 5×10^6 Jyg mammary cancer cells were transplanted subcutaneously in the dorsal region. After 5 days, an osmotic pressure pump (Alzet) containing HGF/NK4 or saline (control) was implanted beneath the dorsal skin and HGF/NK4 or saline was continuously infused into the neighborhood of the transplanted tumor for 2 weeks. The volume of the tumor graft was serially measured and the number of metastatic foci on the lung surface was counted at 28 days after transplantation.

The time course of tumor volume is shown in Fig. 13 A and the number of metastatic foci on the lung surface at day 28 after transplantation is shown in Table 13 B. It will be apparent that whereas the control tumor perfused with saline showed rapid growth at day 10 after transplantation and onwards, HGF/NK4 suppressed growth of the tumor. Regarding pulmonary metastasis, whereas a large number of pulmonary metastatic foci were observed in the control saline-perfused group, pulmonary metastasis was inhibited in the animals perfused with HGF/NK4.

These results indicated that HGF/NK4 has an action to inhibit the growth and metastasis of Jyg mammary cancer significantly in vivo.

Formulation Example 1

A solution containing the HGF/NK4 prepared in Example 1 (1 mg), mannitol (1 g) and polysolvate 80 (10 mg) in 100 ml of saline was aseptically prepared and distributed into vials, 1 ml per vial. The vial contents were lyophilized and the vials sealed to provide the neovascularization inhibitor of the invention in the form of a lyophilizate.

Formulation Example 2

An aqueous solution containing 1 mg of the HGF/NK4 prepared in Example 1 and 100 mg of human serum albumin was aseptically formulated with 100 ml of 0.02 M phosphate buffer (containing 0.15 M NaCl and 0.01% polysolvate 80; pH 7.4) and the mixture was distributed into vials, 1 ml per vial. The liquid contents of the vials were lyophilized and the vials sealed to provide the neovascularization inhibitor in the form of a lyophilizate.

Formulation Example 3

A solution containing the HGF/NK4 prepared in Example 1 (1 mg), sorbitol (2 g), glycine (2 g) and polysolvate 80 (10 mg) in 100 ml of distilled water

for injection was aseptically prepared and distributed into vials, 1 ml per vial. The contents of the vials were lyophilized and the vials sealed to provide the neovascularization inhibitor in the form of a lyophilizate.

INDUSTRIAL APPLICABILITY

The neovascularization inhibitor of the present invention finds application, based on its neovascularization inhibitory activity, as a prophylactic or therapeutic agent for various diseases associated with abnormal angiopoiesis, such as rheumatoid arthritis, diabetic retinopathy, retinopathy of prematurity, senile macular degeneration, and overcicatrization associated with wound healing.

CLAIMS

1. A neovascularization inhibitor comprising the following polypeptide (a) or (b) as an active ingredient:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

2. (amended) A neovascularization inhibitor comprising the following polypeptide (a) or (b) as an active ingredient:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or

VEGF.

3. (amended) The neovascularization inhibitor according to Claim 1 or 2, wherein said polypeptide has at least one hairpin domain and 4 Kringle domains.

4. (amended) The neovascularization inhibitor according to Claim 1 or 2, wherein said polypeptide is one obtainable by elastase digestion of hepatocyte growth factor.

5. (amended) A neovascularization inhibitor comprising the polypeptide defined by SEQ ID NO:1 and a pharmaceutically acceptable carrier.

6. (amended) A neovascularization inhibitor comprising the polypeptide defined by SEQ ID NO:2 and a pharmaceutically acceptable carrier.

7. (amended) A prophylactic or therapeutic drug for a disease associated with abnormal angiopoiesis which comprises the polypeptide defined in Claim 1 or 2 and a pharmaceutically acceptable carrier.

8. (amended) The prophylactic or therapeutic drug according to Claim 7 wherein said disease associated with abnormal angiopoiesis is selected from the group consisting of rheumatoid arthritis, psoriasis, Osler-Webber syndrome, myocardial

angiopoiesis, telangiectasia, hemophilic joint, angiogenic diseases of the eye, angiofibroma, benign tumors and wound granulation.

9. (amended) A prophylactic or therapeutic drug for a disease arising from overstimulation of endothelial cells which comprises the polypeptide defined in claim 1 or 2 and a pharmaceutically acceptable carrier.

10. (amended) The prophylactic or therapeutic drug according to Claim 9 wherein said disease arising from overstimulation of endothelial cells is selected from the group consisting of enteric adhesion, Crohn's disease, atherosclerosis, scleroderma and overcicatrizization.

11. (amended) A conception-regulating drug comprising the polypeptide defined in Claim 1 or 2 and a pharmaceutically acceptable carrier.

12. (amended) A method of inhibiting neovascularization which comprises administering to a subject a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence $\text{PyrGlu}^{32} \sim \text{Val}^{478}$ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence

53/1

derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

13. (amended) A method of inhibiting neovascularization which comprises administering to a subject a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF.

14. (amended) A method for prophylaxis or therapy of a disease associated with abnormal angiopoiesis which comprises administering a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a

53/2

pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF

to a subject in whom a prophylactic or therapeutic treatment for said disease is indicated.

15. (added) A method for prophylaxis or therapy of a disease associated with abnormal angiopoiesis which comprises administering a neovascularization inhibitor comprising the following polypeptide (a) or (b) and a pharmaceutically acceptable carrier:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or

AW 34

53/3

VEGF

to a subject in whom a prophylactic or therapeutic treatment for said disease is indicated.

16. (added) The method for prophylaxis or therapy according to Claim 14 or 15 wherein said disease is any disease selected from the group consisting of rheumatoid arthritis, psoriasis, Osler-Webber syndrome, myocardial angiopoiesis, telangiectasia, hemophilic joint, angiogenic diseases of the eye, angiofibroma, benign tumors, wound granulation, enteric adhesion, Crohn's disease, atherosclerosis, scleroderma and overcicatrizization.

17. (added) Use of the following polypeptide (a) or (b) for the production of a neovascularization inhibitor:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).

(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF.

18. (added) Use of the following polypeptide

(a) or (b) for the production of a neovascularization inhibitor:

(a) a polypeptide having the amino acid sequence PyrGlu³² ~ Val⁴⁷⁸ of hepatocyte growth factor (HGF).


(b) a polypeptide having an amino acid sequence derived from the amino acid sequence defined in (a) by the deletion, substitution or addition of one or more amino acids, and having antagonistic activity against the c-Met/HGF receptor-mediated action of HGF and inhibitory action against the growth of vascular endothelial cells induced by bFGF and/or VEGF.

ABSTRACT

Novel neovascularization inhibitory factors and neovascularization inhibitors useful in preventing and treating various diseases in association with neovascularization. These neovascularization inhibitors contain as the active ingredient polypeptides with the following definition (a) or (b): (a) a polypeptide having an amino acid sequence of PyrGlu³²-Val⁴⁷⁸ in HGF (hepatocyte growth factor); or (b) a polypeptide having an amino acid sequence derived from the amino acid sequence as defined in (a) by deletion, substitution or addition of one or several amino acids and having an antagonism to the effect of HGF via c-Met-HGF receptor.

FIG. 1

A



B




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7





FIG. 8







*FIG. 9***Control****NK4 20 μ g****× 8**

FIG. 10**Control**

* Immunostaining for endothelial cell-specific von Willebrand factor antibody.

FIG. 11





FIG. 12

A



B





FIG. 13



Declaration and Power of Attorney for Patent Application

特許出願宣言書及び委任状

Japanese Language Declaration

日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者（下記の氏名が一つの場合）もしくは最初かつ共同発明者であると（下記の名称が複数の場合）信じています。

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

NEOVASCULARIZATION INHIBITORS

上記発明の明細書（下記の欄でX印がついていない場合は、本書に添付）は、

the specification of which is attached hereto unless the following box is checked:

☐ ____月 ____日に提出され、米国出願番号または特許協定条約

☒ was filed on April 6, 1999
as United States Application Number or
PCT International Application Number

国際出願番号を _____ とし、

PCT/JP99/01834 and was amended on

（該当する場合） _____ に訂正されました。

February 7, 2000 (if applicable).

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第37編第1条56項に定義されるとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

Japanese Language Declaration

(日本語宣言書)

私は、米国法典第35編第119条(a)-(d)項又は第365条(b)項に基づき下記の、米国以外の国の少なくとも一カ国を指定している特許協力条約第365条(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願についての外国優先権をここに主張するとともに、優先権を主張している本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

Prior Foreign Applications

外国での先行出願

1998-134681	Japan
(Number)	(Country)
(番号)	(国名)
(Number)	(Country)
(番号)	(国名)
(Number)	(Country)
(番号)	(国名)

私は、第35編米国法典119条(e)項に基づいて下記の米国特許出願規定に記載された権利をここに主張致します。

(Application No.)	(Filing Date)
(出願番号)	(出願日)

私は、下記の米国法典第35編第120条に基づいて下記の米国特許出願に記載された権利、又は米国を指定している特許協力条約第365条(c)に基づく権利をここに主張します。又、本出願の各請求範囲の内容が米国法典第35編第112条第1項又は特許協力条約で規定された方法で先行する米国特許出願に開示されていない限り、その先行米国出願書提出日以降で本出願書の日本国内又は特許協力条約国際出願提出日までの期間中に入手された、連邦規則法典第37編第1条第56項で定義された特許資格の有無に関する重要な情報について開示義務があることを認識しています。

(Application No.)	(Filing Date)
(出願番号)	(出願日)
(Application No.)	(Filing Date)
(出願番号)	(出願日)

私は、私自身の知識に基づいて本宣言中で私が行う表明が真実であり、かつ私の入手した情報と私の信ずるところに基づく表明が全て真実であると信じていること、さらに故意になされた虚偽の表明及びそれと同等の行為は米国法典第18編第1001条に基づき、罰金または拘禁、もしくはその両方により処罰されること、そしてそのような故意による虚偽の声明を行えば、出願した、又は既に許可された特許の有効性が失われることを認識し、よってここに上記のごとく宣誓を致します。

I hereby claim foreign priority under Title 35, United States Code, Section 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Priority Not Claimed

優先権主張なし

28/04/1998	<input type="checkbox"/>
(Day/Month/Year Filed)	
(出願年月日)	
(Day/Month/Year Filed)	<input type="checkbox"/>
(出願年月日)	
(Day/Month/Year Filed)	<input type="checkbox"/>
(出願年月日)	

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.)	(Filing Date)
(出願番号)	(出願日)

I hereby claim the benefit of Title 35, United States Code Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose any material information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

(Status: Patented, Pending, Abandoned)
(現況: 特許許可済、係属中、放棄済)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Japanese Language Declaration

(日本語宣言書)

委任状：私は、下記の発明者として、本出願に関する一切の手続きを米国特許商標局に対して遂行する弁理士又は代理人として、下記のことを指名致します。(弁理士、又は代理人の氏名及び登録番号を明記のこと)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith (list name and registration number)

John H. Mion, Reg. No. 18,879; Thomas J. Macpeak, Reg. No. 19,292; Robert J. Seas, Jr., Reg. No. 21,092; Darryl Mexic, Reg. No. 23,063; Robert V. Sloan, Reg. No. 22,775; Peter D. Olexy, Reg. No. 24,513; J. Frank Osha, Reg. No. 24,625; Waddell A. Biggart, Reg. No. 24,861; Louis Gubinsky, Reg. No. 24,835; Neil B. Siegel, Reg. No. 25,200; David J. Cushing, Reg. No. 28,703; John R. Inge, Reg. No. 26,916; Joseph J. Ruch, Jr., Reg. No. 26,577; Sheldon I. Landsman, Reg. No. 25,430; Richard C. Turner, Reg. No. 29,710; Howard L. Bernstein, Reg. No. 25,665; Alan J. Kasper, Reg. No. 25,426; Kenneth J. Burchfiel, Reg. No. 31,333; Gordon Kit, Reg. No. 30,764; Susan J. Mack, Reg. No. 30,951; Frank L. Bernstein, Reg. No. 31,484; Mark Boland, Reg. No. 32,197; William H. Mandir, Reg. No. 32,156; Scott M. Daniels, Reg. No. 32,562; Brian W. Hannon, Reg. No. 32,778; Abraham J. Rosner, Reg. No. 33,276; Bruce E. Kramer, Reg. No. 33,725; Paul F. Neils, Reg. No. 33,102 and Brett S. Sylvester, Reg. No. 32,765

書類送付先:

Send Correspondence to:

SUGHRUE, MION, ZINN, MACPEAK & SEAS, PLLC
2100 Pennsylvania Avenue, N.W., Washington, D.C. 20037-3202

直通電話連絡先: (名称及び電話番号)

Direct Telephone Calls to: (name and telephone number)

(202)293-7060

唯一又は第一発明者名	Full name of sole or first inventor	
	NAKAMURA Toshikazu	
発明者の署名	日付	Inventor's signature Date
		T. Nakamura May 31, 2000
住所	Residence	
	Takatsuki-shi, Osaka 569-1020, Japan	
国籍	Citizenship	
	Japan	
郵便の宛先	Post office address	
	4-1, Takamidai, Takatsuki-shi, Osaka 569-1020, Japan	
第二共同発明者名 (該当する場合)	Full name of second joint inventor, if any	
第二発明者の署名	日付	Second inventor's signature Date
住所	Residence	
国籍	Citizenship	
郵便の宛先	Post office address	

(第三以降の共同発明者についても同様に記載し、署名をする (Supply similar information and signature for third and subsequent joint inventors.))

09674377 103000

09/674377

534 Rec'd PCT/PTG 30 OCT 2000

SEQUENCE LISTING

<110> Nakamura, Toshikazu

<120> Neovascularization Inhibitors

<130> Q61434

<140>

<141>

<150> PCT/JP99/01834

<151> 1999-04-06

<150> JP 1998/134681

<151> 1998-04-28

<160> 2

<170> PatentIn Ver. 2.1

<210> 1

<211> 447

<212> PRT

<213> Homo sapiens

<220>

<221> MOD_RES

<222> (1)

<223> pyroglutamate

<220>

<221> CHAIN

<222> (1)..(447)

<223> N-terminal region of alpha-chain in HGF

.. (PyrGlu32-Val478/HGF)

<300>

<301> Nakamura, Toshikazu

<303> Nature

<304> 342

<306> 440-443

<307> 1989

<400> 1

Xaa Arg Lys Arg Arg Asn Thr Ile His Glu Phe Lys Lys Ser Ala Lys

1

5

10

15

Thr Thr Leu Ile Lys Ile Asp Pro Ala Leu Lys Ile Lys Thr Lys Lys
20 25 30

Val Asn Thr Ala Asp Gln Cys Ala Asn Arg Cys Thr Arg Asn Lys Gly
35 40 45

Leu Pro Phe Thr Cys Lys Ala Phe Val Phe Asp Lys Ala Arg Lys Gln
50 55 60

Cys Leu Trp Phe Pro Phe Asn Ser Met Ser Ser Gly Val Lys Lys Glu
65 70 75 80

Phe Gly His Glu Phe Asp Leu Tyr Glu Asn Lys Asp Tyr Ile Arg Asn
85 90 95

Cys Ile Ile Gly Lys Gly Arg Ser Tyr Lys Gly Thr Val Ser Ile Thr
100 105 110

Lys Ser Gly Ile Lys Cys Gln Pro Trp Ser Ser Met Ile Pro His Glu
115 120 125

His Ser Phe Leu Pro Ser Ser Tyr Arg Gly Lys Asp Leu Gln Glu Asn
130 135 140

Tyr Cys Arg Asn Pro Arg Gly Glu Glu Gly Gly Pro Trp Cys Phe Thr
145 150 155 160

Ser Asn Pro Glu Val Arg Tyr Glu Val Cys Asp Ile Pro Gln Cys Ser
165 170 175

Glu Val Glu Cys Met Thr Cys Asn Gly Glu Ser Tyr Arg Gly Leu Met
180 185 190

Asp His Thr Glu Ser Gly Lys Ile Cys Gln Arg Trp Asp His Gln Thr
195 200 205

Pro His Arg His Lys Phe Leu Pro Glu Arg Tyr Pro Asp Lys Gly Phe
210 215 220

Asp Asp Asn Tyr Cys Arg Asn Pro Asp Gly Gln Pro Arg Pro Trp Cys
225 230 235 240

Tyr Thr Leu Asp Pro His Thr Arg Trp Glu Tyr Cys Ala Ile Lys Thr
245 250 255

Cys Ala Asp Asn Thr Met Asn Asp Thr Asp Val Pro Leu Glu Thr Thr
260 265 270

Glu Cys Ile Gln Gly Gln Gly Glu Gly Tyr Arg Gly Thr Val Asn Thr
 275 280 285

Ile Trp Asn Gly Ile Pro Cys Gln Arg Trp Asp Ser Gln Tyr Pro His
 290 295 300

Glu His Asp Met Thr Pro Glu Asn Phe Lys Cys Lys Asp Leu Arg Glu
 305 310 315 320

Asn Tyr Cys Arg Asn Pro Asp Gly Ser Glu Ser Pro Trp Cys Phe Thr
 325 330 335

Thr Asp Pro Asn Ile Arg Val Gly Tyr Cys Ser Gln Ile Pro Asn Cys
 340 345 350

Asp Met Ser His Gly Gln Asp Cys Tyr Arg Gly Asn Gly Lys Asn Tyr
 355 360 365

Met Gly Asn Leu Ser Gln Thr Arg Ser Gly Leu Thr Cys Ser Met Trp
 370 375 380

Asp Lys Asn Met Glu Asp Leu His Arg His Ile Phe Trp Glu Pro Asp
 385 390 395 400

Ala Ser Lys Leu Asn Glu Asn Tyr Cys Arg Asn Pro Asp Asp Asp Ala
 405 410 415

His Gly Pro Trp Cys Tyr Thr Gly Asn Pro Leu Ile Pro Trp Asp Tyr
 420 425 430

Cys Pro Ile Ser Arg Cys Glu Gly Asp Thr Thr Pro Thr Ile Val
 435 440 445

<210> 2

<211> 442

<212> PRT

<213> Homo sapiens

<220>

<221> CHAIN

<222> (1)..(442)

<223> N-terminal region of alpha-chain in HGF
 (PyrGlu32-Val478/HGF)

<220>

<221> MOD_RES

<222> (130)..(131)

<223> deletion of 5 amino acids

<400> 2

Xaa	Arg	Lys	Arg	Arg	Asn	Thr	Ile	His	Glu	Phe	Lys	Lys	Ser	Ala	Lys	1	5	10	15
Thr	Thr	Leu	Ile	Lys	Ile	Asp	Pro	Ala	Leu	Lys	Ile	Lys	Thr	Lys	Lys	20	25	30	
Val	Asn	Thr	Ala	Asp	Gln	Cys	Ala	Asn	Arg	Cys	Thr	Arg	Asn	Lys	Gly	35	40	45	
Leu	Pro	Phe	Thr	Cys	Lys	Ala	Phe	Val	Phe	Asp	Lys	Ala	Arg	Lys	Gln	50	55	60	
Cys	Leu	Trp	Phe	Pro	Phe	Asn	Ser	Met	Ser	Ser	Gly	Val	Lys	Lys	Glu	65	70	75	80
Phe	Gly	His	Glu	Phe	Asp	Leu	Tyr	Glu	Asn	Lys	Asp	Tyr	Ile	Arg	Asn	85	90	95	
Cys	Ile	Ile	Gly	Lys	Gly	Arg	Ser	Tyr	Lys	Gly	Thr	Val	Ser	Ile	Thr	100	105	110	
Lys	Ser	Gly	Ile	Lys	Cys	Gln	Pro	Trp	Ser	Ser	Met	Ile	Pro	His	Glu	115	120	125	
His	Ser	Tyr	Arg	Gly	Lys	Asp	Leu	Gln	Glu	Asn	Tyr	Cys	Arg	Asn	Pro	130	135	140	
Arg	Gly	Glu	Glu	Gly	Gly	Pro	Trp	Cys	Phe	Thr	Ser	Asn	Pro	Glu	Val	145	150	155	160
Arg	Tyr	Glu	Val	Cys	Asp	Ile	Pro	Gln	Cys	Ser	Glu	Val	Glu	Cys	Met	165	170	175	
Thr	Cys	Asn	Gly	Glu	Ser	Tyr	Arg	Gly	Leu	Met	Asp	His	Thr	Glu	Ser	180	185	190	
Gly	Lys	Ile	Cys	Gln	Arg	Trp	Asp	His	Gln	Thr	Pro	His	Arg	His	Lys	195	200	205	
Phe	Leu	Pro	Glu	Arg	Tyr	Pro	Asp	Lys	Gly	Phe	Asp	Asp	Asn	Tyr	Cys	210	215	220	
Arg	Asn	Pro	Asp	Gly	Gln	Pro	Arg	Pro	Trp	Cys	Tyr	Thr	Leu	Asp	Pro	225	230	235	240

His Thr Arg Trp Glu Tyr Cys Ala Ile Lys Thr Cys Ala Asp Asn Thr
 245 250 255

Met Asn Asp Thr Asp Val Pro Leu Glu Thr Thr Glu Cys Ile Gln Gly
 260 265 270

Gln Gly Glu Gly Tyr Arg Gly Thr Val Asn Thr Ile Trp Asn Gly Ile
 275 280 285

Pro Cys Gln Arg Trp Asp Ser Gln Tyr Pro His Glu His Asp Met Thr
 290 295 300

Pro Glu Asn Phe Lys Cys Lys Asp Leu Arg Glu Asn Tyr Cys Arg Asn
 305 310 315 320

Pro Asp Gly Ser Glu Ser Pro Trp Cys Phe Thr Thr Asp Pro Asn Ile
 325 330 335

Arg Val Gly Tyr Cys Ser Gln Ile Pro Asn Cys Asp Met Ser His Gly
 340 345 350

Gln Asp Cys Tyr Arg Gly Asn Gly Lys Asn Tyr Met Gly Asn Leu Ser
 355 360 365

Gln Thr Arg Ser Gly Leu Thr Cys Ser Met Trp Asp Lys Asn Met Glu
 370 375 380

Asp Leu His Arg His Ile Phe Trp Glu Pro Asp Ala Ser Lys Leu Asn
 385 390 395 400

Glu Asn Tyr Cys Arg Asn Pro Asp Asp Ala His Gly Pro Trp Cys
 405 410 415

Tyr Thr Gly Asn Pro Leu Ile Pro Trp Asp Tyr Cys Pro Ile Ser Arg
 420 425 430

Cys Glu Gly Asp Thr Thr Pro Thr Ile Val
 435 440

09674377 09/674377

1/8

534 Rec'd PCT/PTC 30 OCT 2000

SEQUENCE LISTING

<110> Nakamura, Toshikazu

<120> Inhibitor of Vascularization

<130> P99-10

<140>

<141>

<150> JP P1998/134681

<151> 1998-04-28

<160> 2

<170> PatentIn Ver. 2.0

<210> 1

<211> 447

<212> PRT

<213> Human

<220>

<221> MOD_RES

<222> (1)

<223> pyroglutamate

<220>

<221> CHAIN

2/8

<222> (1).. (447)

<223> N-terminal region of alpha-chain in HGF
(PyrGlu32-Val478/HGF)

<300>

<301> Nakamura, Tshikazu

<303> Nature

<304> 342

<306> 440-443

<307> 1989

<400> 1

Xaa Arg Lys Arg Arg Asn Thr Ile His Glu Phe Lys Lys Ser Ala Lys

1

5

10

15

Thr Thr Leu Ile Lys Ile Asp Pro Ala Leu Lys Ile Lys Thr Lys Lys

20

25

30

Val Asn Thr Ala Asp Gln Cys Ala Asn Arg Cys Thr Arg Asn Lys Gly

35

40

45

Leu Pro Phe Thr Cys Lys Ala Phe Val Phe Asp Lys Ala Arg Lys Gln

50

55

60

Cys Leu Trp Phe Pro Phe Asn Ser Met Ser Ser Gly Val Lys Lys Glu

65

70

75

80

Phe Gly His Glu Phe Asp Leu Tyr Glu Asn Lys Asp Tyr Ile Arg Asn

85

90

95

3/8

Cys Ile Ile Gly Lys Gly Arg Ser Tyr Lys Gly Thr Val Ser Ile Thr
100 105 110

Lys Ser Gly Ile Lys Cys Gln Pro Trp Ser Ser Met Ile Pro His Glu
115 120 125

His Ser Phe Leu Pro Ser Ser Tyr Arg Gly Lys Asp Leu Gln Glu Asn
130 135 140

Tyr Cys Arg Asn Pro Arg Gly Glu Glu Gly Gly Pro Trp Cys Phe Thr
145 150 155 160

Ser Asn Pro Glu Val Arg Tyr Glu Val Cys Asp Ile Pro Gln Cys Ser
165 170 175

Glu Val Glu Cys Met Thr Cys Asn Gly Glu Ser Tyr Arg Gly Leu Met
180 185 190

Asp His Thr Glu Ser Gly Lys Ile Cys Gln Arg Trp Asp His Gln Thr
195 200 205

Pro His Arg His Lys Phe Leu Pro Glu Arg Tyr Pro Asp Lys Gly Phe
210 215 220

Asp Asp Asn Tyr Cys Arg Asn Pro Asp Gly Gln Pro Arg Pro Trp Cys
225 230 235 240

Tyr Thr Leu Asp Pro His Thr Arg Trp Glu Tyr Cys Ala Ile Lys Thr
245 250 255

4/8

Cys Ala Asp Asn Thr Met Asn Asp Thr Asp Val Pro Leu Glu Thr Thr
260 265 270

Glu Cys Ile Gln Gly Gln Gly Glu Gly Tyr Arg Gly Thr Val Asn Thr
275 280 285

Ile Trp Asn Gly Ile Pro Cys Gln Arg Trp Asp Ser Gln Tyr Pro His
290 295 300

Glu His Asp Met Thr Pro Glu Asn Phe Lys Cys Lys Asp Leu Arg Glu
305 310 315 320

Asn Tyr Cys Arg Asn Pro Asp Gly Ser Glu Ser Pro Trp Cys Phe Thr
325 330 335

Thr Asp Pro Asn Ile Arg Val Gly Tyr Cys Ser Gln Ile Pro Asn Cys
340 345 350

Asp Met Ser His Gly Gln Asp Cys Tyr Arg Gly Asn Gly Lys Asn Tyr
355 360 365

Met Gly Asn Leu Ser Gln Thr Arg Ser Gly Leu Thr Cys Ser Met Trp
370 375 380

Asp Lys Asn Met Glu Asp Leu His Arg His Ile Phe Trp Glu Pro Asp
385 390 395 400

Ala Ser Lys Leu Asn Glu Asn Tyr Cys Arg Asn Pro Asp Asp Asp Ala
405 410 415

5/8

His Gly Pro Trp Cys Tyr Thr Gly Asn Pro Leu Ile Pro Trp Asp Tyr

420

425

430

Cys Pro Ile Ser Arg Cys Glu Gly Asp Thr Thr Pro Thr Ile Val

435

440

445

<210> 2

<211> 442

<212> PRT

<213> Human

<220>

<221> CHAIN

<222> (1).. (442)

<223> N-terminal region of alpha-chain in HGF
(pyrGlu32-Val478/HGF)

<220>

<221> MOD_RES

<222> (130).. (131)

<223> deletion of 5 amino acids

<400> 2

Xaa Arg Lys Arg Arg Asn Thr Ile His Glu Phe Lys Lys Ser Ala Lys

1

5

10

15

Thr Thr Leu Ile Lys Ile Asp Pro Ala Leu Lys Ile Lys Thr Lys Lys

20

25

30

6/8

Val Asn Thr Ala Asp Gln Cys Ala Asn Arg Cys Thr Arg Asn Lys Gly

35

40

45

Leu Pro Phe Thr Cys Lys Ala Phe Val Phe Asp Lys Ala Arg Lys Gln

50

55

60

Cys Leu Trp Phe Pro Phe Asn Ser Met Ser Ser Gly Val Lys Lys Glu

65

70

75

80

Phe Gly His Glu Phe Asp Leu Tyr Glu Asn Lys Asp Tyr Ile Arg Asn

85

90

95

Cys Ile Ile Gly Lys Gly Arg Ser Tyr Lys Gly Thr Val Ser Ile Thr

100

105

110

Lys Ser Gly Ile Lys Cys Gln Pro Trp Ser Ser Met Ile Pro His Glu

115

120

125

His Ser Tyr Arg Gly Lys Asp Leu Gln Glu Asn Tyr Cys Arg Asn Pro

130

135

140

Arg Gly Glu Glu Gly Gly Pro Trp Cys Phe Thr Ser Asn Pro Glu Val

145

150

155

160

Arg Tyr Glu Val Cys Asp Ile Pro Gln Cys Ser Glu Val Glu Cys Met

165

170

175

Thr Cys Asn Gly Glu Ser Tyr Arg Gly Leu Met Asp His Thr Glu Ser

180

185

190

7/8

Gly Lys Ile Cys Gln Arg Trp Asp His Gln Thr Pro His Arg His Lys

195

200

205

Phe Leu Pro Glu Arg Tyr Pro Asp Lys Gly Phe Asp Asp Asn Tyr Cys

210

215

220

Arg Asn Pro Asp Gly Gln Pro Arg Pro Trp Cys Tyr Thr Leu Asp Pro

225

230

235

240

His Thr Arg Trp Glu Tyr Cys Ala Ile Lys Thr Cys Ala Asp Asn Thr

245

250

255

Met Asn Asp Thr Asp Val Pro Leu Glu Thr Thr Glu Cys Ile Gln Gly

260

265

270

Gln Gly Glu Gly Tyr Arg Gly Thr Val Asn Thr Ile Trp Asn Gly Ile

275

280

285

Pro Cys Gln Arg Trp Asp Ser Gln Tyr Pro His Glu His Asp Met Thr

290

295

300

Pro Glu Asn Phe Lys Cys Lys Asp Leu Arg Glu Asn Tyr Cys Arg Asn

305

310

315

320

Pro Asp Gly Ser Glu Ser Pro Trp Cys Phe Thr Thr Asp Pro Asn Ile

325

330

335

Arg Val Gly Tyr Cys Ser Gln Ile Pro Asn Cys Asp Met Ser His Gly

340

345

350

8/8

Gln Asp Cys Tyr Arg Gly Asn Gly Lys Asn Tyr Met Gly Asn Leu Ser
355 360 365

Gln Thr Arg Ser Gly Leu Thr Cys Ser Met Trp Asp Lys Asn Met Glu
370 375 380

Asp Leu His Arg His Ile Phe Trp Glu Pro Asp Ala Ser Lys Leu Asn
385 390 395 400

Glu Asn Tyr Cys Arg Asn Pro Asp Asp Asp Ala His Gly Pro Trp Cys
405 410 415

Tyr Thr Gly Asn Pro Leu Ile Pro Trp Asp Tyr Cys Pro Ile Ser Arg
420 425 430

Cys Glu Gly Asp Thr Thr Pro Thr Ile Val
435 440